

Flavonoid Glycosides from *Hosta longipes*, Their Inhibition on NO Production, and Nerve Growth Factor Inductive Effects

Chung Sub Kim,^a Oh Wook Kwon,^b Sun Yeou Kim^{c,d} and Kang Ro Lee^{*,a}

^aNatural Products Laboratory, School of Pharmacy, Sungkyunkwan University, 440-746 Suwon, Republic of Korea

^bGraduate School of East-West Medical Science, Kyung Hee University Global Campus, 446-701 Yongin, Republic of Korea

College of Pharmacy, Gachon University, 406-799 Incheon, Republic of Korea

^dGachon Medical Research Institute, Gil Medical Center, 405-760 Incheon, Republic of Korea




Figure S1. HR-ESI-MS spectrum of 1.




Figure S2. ¹H NMR spectrum of 1 (CD₃OD, 900 MHz).




Figure S3. ¹³C NMR spectrum of 1 (CD₃OD, 225 MHz).




Figure S4. ¹H-¹H COSY spectrum of 1.




Figure S5. HMQC spectrum of 1.




Figure S6. HMBC spectrum of 1.




Figure S7. HR-FAB-MS spectrum of 2.




Figure S8. ¹H NMR spectrum of 2 (CD₃OD, 500 MHz).




Figure S9. ¹³C NMR spectrum of 2 (CD₃OD, 125 MHz).




Figure S10. ¹H-¹H COSY spectrum of 2.





Figure S11. HMQC spectrum of 2.








Figure S13. ¹H NMR spectrum of 5 (DMSO-d₆, 500 MHz).