

SYNTHESES, CHARACTERIZATION AND ANTIFUNGAL ACTIVITY OF TRIS(1,10-PHENANTHROLINE) IRON(II) BIS(N-R-SULFONYLDITHIOCARBIMATE)ZINCATE(II)

Fernanda C. Bottega, Marcelo R. L. Oliveira*, Camila V. Garcia, Daniele C. Menezes e Mayura M. M. Rubinger

Departamento de Química, Universidade Federal de Viçosa, 36570-000 Viçosa - MG, Brasil

Laercio Zambolim

Departamento de Fitopatologia, Universidade Federal de Viçosa, 36570-000 Viçosa - MG, Brasil




Figure 1S. Infrared spectrum of compound 1 (CsI pellet)




Figure 2S. Infrared spectrum of compound 2 (CsI pellet)

*e-mail: marcelor@ufv.br




Figure 3S. Infrared spectrum of compound 3 (CsI pellet)




Figure 4S. Infrared spectrum of compound 4 (CsI pellet)




Figure 5S. ^1H NMR spectrum of compound 2 (DMSO-d_6)




Figure 6S. ^{13}C NMR spectrum of compound 2 (DMSO-d_6)




Figure 7S. ^1H NMR spectrum of compound 3 (DMSO-d_6)




Figure 8S. ^{13}C NMR spectrum of compound 3 (DMSO-d_6)




Figure 9S. ^1H NMR spectrum of compound 4 (DMSO-d_6)




Figure 10S. ^{13}C NMR spectrum of compound 4 (DMSO-d_6)