

2.2 Supplementary Table 2S: Signs and Symptoms at Initial Presentation – Survival Analysis

Variables (%)	All patients (n = 137)*	Survivors (n = 129)	Nonsurvivors or with Severe Disability (n = 6)	p-value**
Vomiting	90	87	2	0.1799
Nausea	19	18	1	1
Abdominal Pain	15	15	0	1
Cold Extremities	5	5	0	1
Respiratory Insufficiency	5	5	0	1
Tachypnea	32	29	2	0.6205
Irritability	2	2	0	1
Lethargy/ Drowsiness	22	22	0	0.5889
Salivation	19	19	0	0.5943
Wheezing	1	1	0	1
Diffuse Sweating	52	49	2	1
Sphincter Relaxation	1	1	0	1
Myosis	1	1	0	1
Muscle Spasms	1	1	0	1
Oliguria	1	1	0	1
Altered Level of Consciousness	6	6	0	1
Hypotension	6	5	0	1
Hypertension	7	7	0	1
Acute Pulmonar Edema	15	12	3	0.0183
Bradycardia	25	24	1	1
Tachycardia	27	26	1	1
Palpitations	1	1	0	1

Hypothermia	3	3	0	1
Hyperthermia	1	1	0	1
Cardiorespirator Arrest	1	1	0	1
Rash	1	1	0	1
Hypotonia	1	1	0	1
Dizziness	1	1	0	1
Paresthesia	1	1	0	1

*Clinical presentation data not available for 9 patients, with 6 in the Survivors group.

**Hypothesis tests comparing the groups were conducted by Fisher's exact tests.