

TABLE S3 | Pairwise mean genetic distance values inter and intra *Astyanax* species from the Paraguaçu River basin using COI marker and K2P model. Values are in percentage. **Astyanax* sp. from BOLD System.

	<i>A. brucutu</i>	<i>A. lorien</i>	<i>A. epiagos</i>	<i>A. rupestris</i>	<i>A. aff. rupestris 1</i>	<i>A. aff. rupestris 2</i>	<i>Astyanax</i> sp. (Coité River)
<i>A. brucutu</i>	0.0						
<i>A. lorien</i>	0.3	0.0					
<i>A. epiagos</i>	1.8	1.4	0.0				
<i>A. rupestris</i>	2.7	2.3	2.9	0.001			
<i>A. aff. rupestris 1</i>	2.9	2.5	3.1	1.8	0.0		
<i>A. aff. rupestris 2</i>	2.9	2.5	3.1	1.8	0.7	0.0	
<i>Astyanax</i> sp. (Coité River)*	1.2	0.8	2.0	2.9	2.7	3.1	0.0
<i>Astyanax</i> sp. (Piabinha River)*	2.9	2.5	3.1	1.8	0.7	0.0	3.1



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2023 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

HOW TO CITE THIS ARTICLE

- Silva-Santos R, Machado CB, Zanata AM, Camelier P, Galetti Jr. PM, Freitas PD.

Molecular characterization of *Astyanax* species (Characiformes: Characidae) from the upper Paraguaçu River basin, a hydrographic system with high endemism. Neotrop Ichthyol. 2023; 21(2):e230032. <https://doi.org/10.1590/1982-0224-2023-0032>

